



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

II. De Curvarum Tangentibus è Maximorum ac Minimorum Theoriâ immediatè deductis : Unâ cùm Theorem : quibusdam ad Sectiones conicas pertinentibus, ejusdem calculi auxilio investigatis. Autore Humphrido Dittono.

Tangentium methodum propono, facilem satis ac generalem, imò generalissimam, ut pote curvis omnibus unâ eâdemque opera inservientem.

Neque novam vocare metuo, cùm celebriorum Geometrarum nullus (in quantum unquam scire potui) aliquid hujus generis publici juris fecit. Pauca tantum ejus specimina hîc in medium profero, nire enim tum clarâ ac apertâ exemplorum multitudine non indigetimus !

Sit Curva $A G H$, cujus vertex A , axis $A K$, ordinatim applicata $F D$ centrumque (siquod habet) punctum K . Sumpto puncto L in Axe sit $A L = n$, $A D = x$, $F D = y$, $F L = z$; quarum quantitatum, tres posteriores sunt fluentes, prior verò n permanens ac stabilis, hæc enim una eademque prioribus variis semper respondet. Ex Triangulo Rectangulo $F D L$, hanc habemus Equationem, $z z = y \dot{y} + n n - 2 n x + x x$; determinandoque z ad extremum, oritur $2 y \dot{y} - 2 n \dot{x} + 2 x \dot{x} = 0$; unde interpretando $2 y \dot{y}$ secundum propriam Curvæ naturam, relinquetur quantitas n exposita in terminis etiam Curvæ propriis.

Cùm verò z hoc modò ad valorem extremum determinatam habeamus; hoc est, linea $F L$ omnium quæ a puncto L ad Curvam duci possunt vel maxima vel mi-

X x x x x x

nima

nima sit, indeque ad Curvam in puncto F normalis; ipsam D L esse subnormalem patet, ex quâ subtangens nullo negotio eruitur.

In exemplum producatür primò Parabola Apolloniana, quàm curvam hîc delineatam esse supponemus.

Habemus ergo $2 y \dot{y} = r \dot{x}$ (posito Parametro $= r$)

unde $r x - 2 n x + 2 x x = 0$, & $n = \frac{r}{2} + x$, ergo

DL subnormalis $= \frac{1}{2} r$. (Cujus Theorematis sensus hîc est, viz. Si ultra terminum D abscissæ A D, designetur D L semiparametro equalis, atque à puncto L producatür L F recta ad punctum F; recta sic ducta Parabolæ in puncto F normalis erit, & omnium quæ à puncto L, ad Curvam duci possunt minima. Dico minimam; alicui enim curvæ naturam ac indolem scienti, apparet Maximam esse non posse (idquod in sequentibus notatum velim) sed necessario est vel maxima vel minima, ideoque posterior.) Hæcque pars prior est Theor. 5, Lib. 7. Conicor. Præclarissimi de La Hire.

Ducatur ordinata E B, junganturque puncta E, L; sit intercepta B D = f, unde A B = x - f, &

$B L = \frac{r}{2} + f$. Jam $L E^2 = \frac{r r}{4} + r x + f f$, &

$F L^2 = \frac{r r}{4} + r x + f f$, & $F L^2 = \frac{r r}{4} + r x$, ergo

$L E^2 - \frac{4}{4} F L^2 = B D^2$; quæ pars posterior est Theorem. 5. ejusdem Lib. Conicor.

Quò propriùs punctum F in quò curvam normalis secat, puncto A sive vertici admovetur; eò propriùs etiam punctum L eidem venit. Ergo quando F cùm A coin-

A coincidit, & sic evanescit ordinata F D, tunc ipsa Minima jacet in Axe A K, & semiparametri quantitatem adequabit. Hoc est in illo casu $n = \frac{1}{2} r$ tantum; in nihilum abeunte x abscissâ ad ordinatam evanescentem pertinenta. Si ergo $AL = n = \frac{1}{2} r$, sumpto puncto D inter A & L, fiat $AD = x$; tum oritur

$$FL^2 = \frac{r^2}{2} + x x, \text{ ergo } FL^2 - AL^2 = x x,$$

hoc est $FL^4 - AL^4 = AD^4$ semper. Eiusdemque tenoris est Theor. 2. Lib. 7. Conicor. de La Hire.

Secundò sit curva quædam ordinis Parabolici superioris, cujus æquatio $r^{\frac{p-q}{p}} x^{\frac{q}{p}} = y^{\frac{p}{p}}$.

$$\text{Tum } y y = r^{\frac{2p-2q}{p}} x^{\frac{2q}{p}}, \text{ adeoque}$$

$$2 y y = \frac{2q}{p} r^{\frac{2p-2q}{p}} x^{\frac{2q-p}{p}} x; \text{ substituendoque hunc va-}$$

lorem loco $2 y y$ in æquatione generali determinante z ad

$$\text{tremum, habemus inde } n = \frac{q}{p} r^{\frac{2p-2q}{p}} x^{\frac{2q-p}{p}} + x; \text{ \&}$$

$$\text{propterea subnormalis } DL = \frac{q}{p} r^{\frac{2p-2q}{p}} x^{\frac{2q-p}{p}}$$

Hoc verò singulis hiscè curvis facillimè applicatur , si indices p & q secundum unius cujusque naturam ac genium debito modo exponantur.

Supponetur tertio Curvam esse Ellipsiu cujus $\frac{1}{2}$ Axis Mador $A K$; ex cujus etiam equatione consequitur

$$2 y \dot{y} = r \dot{x} - \frac{2 r x \dot{x}}{q} \quad \text{Unde provenit}$$

$$r \dot{x} - \frac{2 r x \dot{x}}{q} - 2 n \dot{x} + 2 x \dot{x} = 0, \text{ \&}$$

$$n = \frac{r}{2} + \frac{x}{q} - r x, \text{ ac propterea } r - \frac{r}{2} - \frac{r x}{q}$$

equalis. Sivero ellipseos loco substitueretur Circulus, equationem eodem modo tractando, inveniemus $DL = r - x$, posito r Circuli Radio æquali.

Sed ad Ellipsiu revertendum, cujus alia proprietas ex hoc fonte deducenda est, prout in Parabolâ factum.

$$\text{Sit } BD = f, \text{ unde } AB = x - f. \text{ Habemus } LE^q (= LB^q + EB^q) = \frac{r r}{4} - \frac{r r x}{q} + \frac{r r x x}{q q} + f f + r x -$$

$$\frac{r x x}{q} - \frac{r f f}{q} : \& FL^q (= FD^q + LD^q) = r x - \frac{r x x}{q} + \frac{r r}{4} - \frac{r r x}{q} + \frac{r r x x}{q q} : \text{ Ergo, } LE^q - LF^q = f f - \frac{r f f}{q} ;$$

Hoc verò est Theor. 6. Lib. 7. Conic. de La Hire.

Postulat enim Geometra ille sublimis, ut sit $q. r.$ $q - x : L D$, cujus valor est $r - \frac{r x}{q}$ prout supra in-

$$\frac{r}{2} - \frac{r x}{q}$$

ventum ;

ventum ; ideoque quarta proportionalis est tribus au-
te positis : Hoc verò ei concesso , LF esse minimam
omnium rectarum quæ à puncto L ad Ellipsiu duci
possunt evidenter demonstrat. Preterea quoniam est
 $q : q - r :: f : f - fr$, Ergo $\frac{ff - rff}{q}$

sive $f \times f - fr$, idem est quòd rectangulum apud D
 $\frac{q}{q}$

DeLa Hire exemplar vocatum : Hoc verò exemplar se-
cundum ejus definitionem, est Rectangulum simile Rect-
angulo , differentiam inter Quadratum Axis Transver-
si & Figuram constituenti (hoc est Rectangulo $qq - qr$)
& preterea ad Rectam BD sive f applicatum. Et
quòd Rectangulum $ff - rff$ omnes hæcæ conditiones
possideat, luce Meridianâ ^qClarius est.

Notetur , ex valore quantitatis n supra invento ,
planè consequi $n \approx \frac{r}{2}$. Nam $n = r \div x \div r x$, ergo

$q n \div r x = \frac{qr}{2} \div qx$, sed (propter $q \approx r$) $qx \approx rx$,

ergo, $q n \approx \frac{qr}{2}$, & $n \approx \frac{r}{2}$.

Quando (ut in Parabola modò observatum) pun-
ctum F in A verticem incidit , ipsa Minima in Axe de-
signatur ; & propter evanescentem x , habemus
 $n = r$: Assumptoque quovis puucto D inter A & L ,
si $AD =$ alicui x , comparando emergit $FL^q - AL^q =$
 $xx - \frac{rxx}{q}$; quòd ipsum est Theor. 3. Lib. 7. Conic.
 $\frac{q}{q}$ D. La

D. La Hire. Quoniam enim est $q : q - r :: x : x - r x$, patet $\frac{q}{q} = \frac{xx - rxx}{q}$ esse exemplar, sed ap-

plicatum ad abscissam x ; & preterea hoc esse mensuram adequatam defectûs, quadrati Minimæ à quadrato cujus vis rectæ alterius, ab eodem puncto ad curvam protensæ; hæcque demonstrat ille loco citato.

Theoremata verò ad Axem minorem sive conjugatum ellipsos spectantia (hactenus enim majore sive Transverso usi fuimus) eodem planè modo determinantur. Sit jam $A K$ Axis Minoris $= \frac{c}{2}$, Parameter $= R$;

punctum L jam ultra centrum, ad alteras partes $G K$ collocari supponitur. Operando ut priùs, invenietur $A L$ sive $n = R + x - R x$, & subnormalis $D L = R - R x$;

Hoc est $c : R :: \frac{c}{2} - x : R - R x$, adeoque ducta $F L$

omnium quæ à puncto L ad ellipsu duci possunt Maxima, & $L F^2 - L E^2 = R f f - f f =$ Rectangulo exemplar ad $\frac{c}{c}$

$B D$ (sive f) applicato. Quòd verò hoc sit exemplar, patet, est enim $c : R - c :: f : \frac{R f - f}{c}$, adeoque ex defi-

nitione, $\frac{R f - f}{c} = f x f =$ Exemplari. Hoc verò Theor. est

7 Lib. 7 Conicor. De La Hire.

Iterum;

Iterum ; Puncto F cùm A coincidente ; propter evanescentem x evanescentis tunc temporis ordinatæ, re-

linquitur $n = \frac{R}{2}$, & AL omnium quæ à puncto L

ad Ellipsin duci possunt Maxima, & $AL^2 - FL^2 = Rxx - xx =$ Exemplari ad AD sive x applicato ;

$\frac{c}{c}$ eodemque modo sonat. Theor. 4. Lib. prædicti Conicorum.

Observandum verò ad casum precedentem (quòd priùs ergo notari debuit) ubi invenimus

$n = \frac{R}{2} + x - \frac{Rx}{c}$, quòd $n \simeq \frac{R}{2}$; nam $c n + Rx =$

$\frac{Rc}{2} + cx$, & propter $R \simeq c$, adeoque $Rx \simeq cx$, re-

linquitur $c n \simeq \frac{Rc}{2}$, & $n \simeq \frac{R}{2}$.

Jam verò ut res in Ellipsi peracta est, sic eodem prorsus modo in Hyperbola peragenda foret, Minimæque in hac curvâ lineæ determinandæ : sed talis inter hascæ curvas connectio, tam facilisque ab unâ ad alteram transitus, ut vel Tyronibus ipsis labor inanis videatur. Nil aliud restat, v. gr.

ad subnormalem determinandam, quàm ut signum $-$ in $+$ mutetur. Nam cùm in Hyperbolâ sit

$$2yy = \frac{rx}{q} + 2rxx, \quad \& \quad n = \frac{r}{2} + x + \frac{rx}{q}$$

(exquatione

generali) manet $DL = \frac{r}{2} + \frac{rx}{q}$.

Concipietur Quarto Curvam M S N (in altera Fig. Fig. parte delin. Esse unam ex Hyperboloidibus, cujus Asymptoti A K, K H, rectamque S R ad Asymptoton K H ordinatam, S R sit = y, S P = z, K R = x, K P = n, quæ hîc necessariò minor erit quàm x, ut consideranti patet. Equatio curvæ propria est $y^p x^q = r^q s^p$ cujus loco (propte r & s quantitates determinatas) scribi possit $y^p = x^{-q}$. adeoque

$$y = x^{-\frac{2q}{p}}, \quad \& \quad 2 y \dot{y} = -\frac{2q}{p} \frac{\dot{x}}{x} x^{-\frac{2q}{p}}; \quad \text{hinc cum}$$

$$z z = y y + x x - 2 n x + n n, \quad \text{pro extremo habemus}$$

$$2 y \dot{y} + 2 x \dot{x} - 2 n \dot{x} = 0, \quad \text{hoc est } -\frac{2q}{p} \frac{\dot{x}}{x} x^{-\frac{2q}{p}} + 2 x \dot{x} - 2 n \dot{x} = 0,$$

$$+ 2 x \dot{x} = 2 n \dot{x}, \quad \& \quad n = x - \frac{q}{p} x^{-\frac{2q}{p}}$$

$$\text{adeoque subnormalis P R } (= x - n) = \frac{q}{p} x^{-\frac{2q}{p}}$$

Curvam jam A F G (ultimo loco) Cycloiden primariam concipiamus; sitque r Radius, c Arcus & y ordinata Circuli genitoris, cujus Diameter per A K representatur centrumque inter L & K positum. Tum vocatâ F D cycloidis ordinatâ a, cæterisque ut priùs; curvæ equatio est $a a = y y + 2 c y + c c$, adeoque $z z (= a a + n n - 2 n x + x x) = y y + 2 c y + c c + n n - 2 n x + x x$, & (z ad extremum determinatâ)

$$\text{natâ) } 2y\dot{y} + 2c\dot{y} + 2y\dot{c} + 2cc - 2n\dot{x} + 2x\dot{x} = 0.$$

Est verò $\dot{y} = \frac{r\dot{x} - x\dot{r}}{y}$, & $\dot{c} = \frac{r\dot{x}}{y}$, ergo hos valores sub-

stituendo, ac equationem debite reducendo, habemus
 $2r - \frac{2rc}{y} - 2xc + 2r + \frac{2cr}{y} = 2n - 2x$; ac

propterea $2r - x + \frac{2rc - xc}{y} = n - x = D.L.$
 subnormali.

Incomparabilis D. Barovius subtangente præcognitâ, ad Maximum & Minimum determinandum utitur; hocque idem post eum fecit D. Neiwentit in suâ infinitorum Analyfi. Cùm verò multis aliis Methodis, in quibus nihil omnino de Curvarum Tactione presupponitur, Maximum ac Minimum inveniri queant, palàm est è Maximis & Minimis ad Tangentes determinandas, tutò ac legitimè procedere posse.

COROLL. I.

EXempla hætenus oblata percurrenti, in singulis vis patebit, quòd $2y\dot{y} - 2n\dot{x} + 2x\dot{x} = 0$, posito nempe loco n in hac equatione, valore ejus secundum curvæ naturam, In Hyperboloidibus ergo

$$Yy\dot{y}yy\dot{y}y \quad ex.$$

$$\text{ex.gr. } \frac{2q \cdot}{p} x x - \frac{2q-p}{p} x x + \frac{2q}{p} x x - \frac{2q-p}{p} x x + 2 x x = 0,$$

quod (ipso oculo judice) manifestum est ; & sic in aliis (sine ullâ demonstratione) veritas faciliè perspicietur.

C O R O L L. II.

EX subnormalium inventione , curvarum ordinatas Maximas & Minimas facile determinabimus. Hâcque in re dico, si subnormalis (pro aliquo curvæ puncto) nihilo ponatur equalis, habemus ordinatam istius curvæ ad extremum determinatam ; & quidam maximam si ad partes curvæ concavas, minimam verò si ad convex applicari intelligatur. Ex. gr. in Circulo (positâ subnormali = 1) est $1 = r - x$; sit $r - x = 0$; ergo $r = x$, ac nide $y = r$, hoc est applicata maxima Ra-

dio equalis. Similiter in Ellipfi, $1 = \frac{r}{2} - \frac{rx}{q}$; sit

$$\frac{r}{2} - \frac{rx}{q} = 0, \text{ tum } r q = 2 r x, \text{ ac } x = \frac{q}{2}, \text{ ergo } y = \frac{r q}{4}$$

= 4tæ parti Figuræ (utivocant) sive femiaxis conjugati quadrato , adeque maxima $y =$ isti semaxi. Nec Methodo dissimili cum aliis curvis operandum foret ; inveniatur subnormalis ex equatione datâ, cûque nihilo equali.

equali posita, ordinatam curvæ maximam vel minimam determinatam habebimus; priorem ad partem curvæ versus axem concavam, posteriorem ad convexam.

POSTSCRIPTUM

Prioribus sequentia hæc (notatu non indigna) adjungi possunt.

Primò æquè facile hâc methodo determinari Tangentem, ad partes curvæ convexas operando, ac ad partes concavas uti priùs. Sit enim $A C$ Tangens verticalis inque eâ ad libitum sumpto puncto C , sit $A C = n$, $CO = z$ (quò etiam charactere omnes lineæ, à puncto C ad curvam convexam $A E G$ ductæ, insigniantur) ergo ductâ MO semper ad $A C$ perpendiculari, erit $CM = n - y$, & cum $OM = x$, erit $z z = n n - 2 n y + y y - x x$, adeoque (pro extremo ipsius z valore) $2 y \dot{y} + 2 x \dot{x} - 2 n \dot{y} = 0$. In quâ equatione si exponatur $2 x x$ secundum curvæ naturam, lineam $C Z$ (quæ hoc loco subnormalis vicem subibit) determinatam dabimus. Res clarior est quàm quæ exemplis Illustrantibus indigeat; quæque jamjam dicta sunt facile hoc opus exculabunt.

Secundò, Sicut Methodo priore, (Curvarum Tangentes invenimus) ipfus lineas LE vel CO à puncto
 $Y y y y y y y 2$ dato

dato vel in Axe vel in Tangente verticali sumpto productas, ad extremum determinando ; sic etiam considerando lineas Q E, &c. à puncto in Axe dato ultra verticem productas, idem (idque Universaliter) perficere possumus. Omnes enim lineæ Q E valoris fluentis sunt ac perpetuò mutabiles, sola verò Tangens Q F (posito quòd Q F curvam tangat) stabilis est ac ad unicum valorem determinata. Hec ergo loco, π extremi Hypothesis non innotemur, sed quantitatem permanentem tantum speculabimur. Assumantur duo puncta Q L, indeque ad idem curvæ punctum E duæ semper lineæ ducantur L E, Q E. Inter punctum F contactus ac verticem, angulus Q E L semper erit obtusus, ad alteras verò partes puncti F acutus erit, supposito (quòd prius monitum) Q F curvam tangere, ac F L ei ad angulos rectos insistere. Sit Q A = p. A L = n. A B = x. B E = y. Q E = z. V E (intercepta inter punctum E & V ubi cadit Q V perpendicularis ab Q in L E productam) = v. Jam propter Triangulum obtusangulum Q E habemus hanc equationem


$$z z = p^2 + 2 p n - y^2 - x^2 + \frac{y^2 + n^2 - 2 n x + x x}{x} \\ x = 2 v, \text{ five loci } \frac{y^2 + n^2 - 2 n x + x x}{x} \text{ scribendo } f, \\ \text{est } z z = p^2 + 2 p n - y^2 - x^2 + 2 n x - 2 f v, \text{ ideoque} \\ 2 z \dot{z} = 2 y \dot{y} - 2 x \dot{x} + 2 n \dot{x} - 2 f \dot{v} - 2 v \dot{f}.$$

Si z jam fiat quantitas stabilis, quò in Casu Q E cùm Q F tangente coincidat, erit tum

$$- 2 y \dot{y} - 2 x \dot{x} + 2 n \dot{x} = 0 \text{ (rectangulo } 2 f v \text{ ejus-} \\ \text{que adeo fluxione penitus evanescente.) Hæc verò} \\ \text{est ipsa æquatio Generalis Methodo superiori determinata,} \\ \text{quæque}$$

(345)

quæque uti-videmus non minus facili ac naturaliter ex hoc suppositi quantitatis stabilis principio, quàm ex illo extremi deducitur.



III. Specimen